

Maximisation Principles and Daisyworld

G.J. Ackland

School of Physics, The University of Edinburgh, Mayfield Road,

Edinburgh EH9 3JZ, UK E-mail: gjackland@ed.ac.uk

(Dated: February 2, 2008)

Abstract

We investigate whether the equilibrium time averaged state of a self-organising system with many internal degrees of freedom, 2D- daisyworld, can be described by optimising a single quantity. Unlike physical systems where a principle of maximum energy production has been observed, Daisyworld follows evolutionary dynamics rather than Hamiltonian dynamics. We find that this is sufficient to invalidate the MEP principle, finding instead a different principle, that the system self-organises to a state which maximises the amount of life.

Keywords: Daisyworld, entropy, feedback, gaia, logistic

I. INTRODUCTION

Thermodynamics provides an excellent macroscopic description of the physics of matter in terms of time-independent averaged quantities such as energy, entropy, free energy. These arise from the underlying microscopic, time-dependent motions of atoms. The two descriptions are unified through statistical mechanics, and therefore contain the same information, but in the absence of atomic-level detail the thermodynamic picture is far more useful for everyday application.

It is interesting to consider whether ecosystems can similarly be described by macroscopic averages rather than details of individual species.

Recent work has suggested that the principle of maximum entropy production (MEP) may be a good way to investigate complex, self organising systems. Lorenz et al (2001,2002) suggest that the atmosphere of Earth and Titan may be in such a state. Dewar (2003) has shown that other known distributions of open, dynamical systems, such as self-organised criticality (Bak et al, 1988), the fluctuation theorem (Crooks, 1999) and Zipf's Law (Zipf 1932). The basis of Dewar's argument is that if one takes an appropriate average over all its internal possibilities[1], consistent with the known fluxes and internal energies, MEP will emerge. Of course, real systems do not actually sample all possibilities, so this only provides a sufficient conditions for MEP, the necessary condition is that the average over possibilities actually observed converges rapidly. This parallels an argument in equilibrium statistical mechanics, where thermodynamic averages are calculated from an ergodic hypothesis (that all microstates contribute according to Boltzmann statistics) to which the chaotic trajectories of real systems rapidly converge.

Boltzmann statistics follow from Hamiltonian dynamics. In the case of ecological systems, the dynamics are driven by evolution, and no proof of convergence of an evolving system to a Boltzmann-type distribution yet exists. It is interesting to ask whether more general principles, such as MEP, can be applied in an ecological context. In this paper we explore a particular ecological model, 2D-Daisyworld, in which the MEP state can be rigorously defined, to see whether the system does, in fact, self-organise thereto.

The 2D-Daisyworld, is based on previous work (Von Bloh et al., 1997,1999, Ackland et al 2003). Life is reduced to a single species 'daisies' with a single property (albedo) upon which selection can act. The local environment is reduced to a single variable temperature.

The planet is warmed by a Sun whose heat production (insolation) increases slowly. Growth is only possible across a narrow range of temperatures (5-40°C) with the mid point 22.5°C being optimal and the growth rate dropping parabolically to zero at the extremes. Local temperature is partly determined by insolation and albedo and partly by diffusive heat flow across the planet. Growth occurs by seeding of bare ground adjacent to existing daisy populations, eliminating the need in the original, zero-dimensional daisyworld of Watson and Lovelock, (1983) to predefine different local environments for different albedo types. Each (asexual) offspring mutates in albedo at random.

The original daisyworld is deterministic, cast in the form of differential equations, and has spawned many variants to incorporate specific evolutionary dynamics or natural selection (Lenton, 1998, Lenton and Lovelock 2000, Lovelock 1998, Robertson and Robinson, 1998, Staley 2002, Sugimoto, 2002). The 2D version is stochastic, with many internal degrees of freedom, allowing an entropy production to be defined by the range of albedo (called biodiversity), as well as through heat flow.

Historically, daisyworlds were designed to illustrate the self-regulating effects of coupling between life and the environment, and their impact on evolution by group selection. This is not the issue of interest here - daisyworld is used as a test case for MEP.

II. METHODS

The temperature field changes as:

$$\begin{aligned}
 C\dot{T}(x, y, t) = & D_T \nabla^2 T(x, y, t) \\
 & - \sigma_B T^4(x, y, t) \\
 & + S_L (1 - A(x, y, t))
 \end{aligned} \tag{1}$$

Where $C = 2500$ is the heat capacity, T the temperature field, $D_T = D/C$ the thermal diffusion constant $\sigma_B = 5.67 \times 10^{-8}$ the Stefan-Boltzmann constant, $S_L = S/432.3$, the insolation normalised to that which gives $T=295.5\text{K}$ on a bare planet and A the albedo field (daisy/bare ground). Each lattice point has an associated temperature and albedo. A ranges from 0 to 1, with 0.5 being the value for bare ground. The daisy field is discrete and evolves stochastically. We use a square lattice with eight neighbouring sites. At each time

step each site is examined and:

(1) If occupied by a daisy, it changes to bare ground with probability $\gamma(T)$ (the "death rate").

(2) If occupied by bare ground, the site is populated with probability $\beta(T)$ (the "growth rate").

The new daisy has the albedo of its parent, from a randomly chosen neighbouring site, with a random fluctuation (mutation) drawn from a uniform distribution between r and $-r$. If the neighbouring site is bare, no growth occurs.

III. MEP OR GAIA?

Two global principles will be advanced, these are not new, but formal definitions for the present system are made.

1/ Maximal Entropy Production (Dewar, 2003) - the system self organises to produce entropy as rapidly as possible

2/ The Gaia principle (Lovelock, 1998) - the system self organises to maximise the amount of life

In previous work on the 2D case (Ackland et al 2003), it was shown that multiple solutions satisfied the Gaia principle. A secondary criterion was found, maximisation of biodiversity entropy[2] $N_A \ln N_A$ where N_A is the number of daisies with albedo A , leading to a mean distribution

$$\langle N_A \rangle = e^{\beta A} \quad (2)$$

with β given by the solution of:

$$\langle A \rangle \int_0^1 \exp(\beta A) dA = \int_0^1 A \exp(\beta A) dA \quad (3)$$

Thus the global state of 2D daisyworld was classified in terms of two quantities, maximised growth (i.e. temperature regulation at the favoured growth temperature) and maximised biodiversity. These are equivalent to satisfying both MEP and Gaia principles simultaneously.

It is possible to make a small change to the 2D daisyworld so that Gaia and MEP cannot be satisfied simultaneously. To do this simply requires making the death rate temperature

dependent, and the temperature for minimum death rate (T_d) different from that for maximum birthrate (T_g). In this case the MEP will predict that the temperature will adjust close to T_g , while the Gaia hypothesis predicts it will adjust close to T_d . [3] The explanation for this is straightforward: MEP predicts that entropy (biodiversity) be produced at the fastest possible rate: maximising the birthrate. By contrast, the Gaia hypothesis is that the total number of daisies be maximised: minimising deathrate. Thus investigating whether the mean temperature tends to T_g or T_d determines which (if either) of the two hypotheses applies. Figure.1 shows the variation of mean temperature and population as a function of insolation: the mean temperature obtained is T_d and the population is $1 - \gamma(T_d)$. Thus we conclude that the dynamics satisfy Gaia rather than MEP.

IV. THERMAL ENTROPY PRODUCTION

The arguments above pertain to biodiversity entropy. With a finite diffusivity, one can also investigate whether the MEP can be applied to entropy production by heatflow in daisyworld. There are conceptual difficulties with defining the system here - the models for MEP on Earth (Ozawa et al, 2003), and Dewar's (2002) formulation assume constant flows of energy into and out of the system; the self-organising albedo of daisyworld is able to regulate these flows. Similarly, the diffusion constant is fixed in 2D-daisyworld, whereas in atmospheric models it is assumed to self adjust. Notwithstanding this, there is a range of temperatures in 2D daisyworld, and heat is transported from hotter to colder regions, so we can define thermal entropy production[4] as

$$dS/dT = \int \frac{dQ}{T} = \sum_i \sum_{k=1}^8 \frac{D_T(T_i - T_k)}{2T_i} \quad (4)$$

Given the constraints on flow, it is not straightforward to determine the maximum possible entropy production. However, after an abrupt change in the external forcing (insolation) the system will be out of equilibrium, and will then evolve towards a steady state. If MEP holds, the state immediately after the change should have lower entropy production than the steady state. Figure 3 shows no sign of such behaviour. Monitoring entropy production also enables us to monitor another property, the net heat flow in and out of the system. In steady state, the mean values of these heat flows should be equal - in 0D daisyworlds solutions initially assumed input and output were in balance at all times - this is not required

for a nonlinear system, but proof of the stability of these solutions came later (Saunders, 1994). In the 2D-daisyworld Figure 2 shows the time variation of flow in and flow out. An interesting feature here is that flow in (basically albedo) fluctuates much faster than flow out (basically temperature), which also has a delayed response of many daisy lifetimes.

Observations of planetary MEP are presumed to indicate that large-scale atmospheric heat transfer mechanisms are self-organising. However, even in the curved-planet version of 2D-daisyworld (Ackland et al, 2003), very strong temperature regulation eliminates large scale temperature variation and hence the driving force for the atmosphere (except in the case of desert formation). Figure 3 shows that the mean heat entropy produced per cell does not tend to a maximum during equilibration after a sharp change in insolation. We thus conclude that no MEP principle applies to heatflow in 2D daisyworld.

We examined whether there was an optimum value for the diffusion constant D : at given S , heat entropy is largely independent of D . This is non-trivial, since it implies that for high D temperature variations are suppressed precisely enough to compensate for the increased diffusion constant. However, it indicates that even if D could be optimised by a hypothetical daisyworld atmosphere, in the presence of strong temperature regulation no detectable maximum entropy producing value of D exists.

V. MEP IN ORIGINAL DAISYWORLD

In Watson and Lovelock's (1983) 0D daisyworld two daisy albedos exist, denoted by suffices b and w , with albedo $A_b = 0.25$ and $A_w = 0.75$ respectively. The relevant equations are:

Change of fractional areas occupied by each daisy type (set to zero at equilibrium)

$$da_i/dt = a_i(x\beta_i - \gamma); \quad (5)$$

Fraction of free surface for growth

$$x = 1 - \sum_i a_i; \quad (6)$$

Growth rates for daisies.

$$\beta_i = \max\{0, 1 - ((22.5 - T_i)/17.5)^2\}; \quad (7)$$

Mean planetary albedo (reflectivity)

$$A_p = x/2 + \sum_i A_i a_i; \quad (8)$$

Local temperatures over daisyfields

$$(T_i + 273)^4 = q(A_p - A_i) + (T_e + 273)^4 \quad (9)$$

One further equation closes the set - this represents the equality of absorbed and emitted heat, As we saw in the 2D case, this equality strictly should be considered as a time average, however if we assume that no variable large heat reservoirs (e.g. icecaps) exist the timescales are such that the following equality holds:

$$(T_e + 273)^4 = LS/\sigma_B(1 - A_p) \quad (10)$$

S being an insolation, L a variable of order unity, and emitted by Stefan-Boltzmann radiation.

The usual conclusion from this model is that the daisies moderate the temperature, however the graph of T vs L (figure 4a): shows that T_e is not optimised to the optimum for growth. In contrast, a plot of x vs L (figure 4b) shows that x is a minimum for all L .

The daisyworld equations can be solved analytically. The solutions fall into four independent classes: no daisies, black-only, white-only, both types. At a given insolation only one “living” solution is stable if the simulation is started with non-zero concentration of both daisy types. Dead and living solutions coexist in regions where the dead planet temperature lies outside the limits for growth (5-40°C). The stable solutions (Fig.4) have been found dynamically, (iterating equations 5-10 to self consistency) by numerous authors. The important point here is that the *stable* solution can be determined by minimisation of x among allowed solutions. Adopting the lowest x is the 0D equivalent to the Gaian principle of maximum life, as found in the 2D case.

As a further test, we introduce a third daisy species (grey) with albedo 0.5, the same as bare ground and therefore with no regulating effect. Equations 5 show that coexistence of three species is impossible, and in fact the stable solutions (Fig.5) traverse regions of black-only, black and grey, grey only, grey and white, white only. In each region unstable equilibria exist for other combinations, and in each case the unstable equilibria have higher

x than the total one. This result is robust against changes to q , A_i and γ (including species dependent γ).

The equilibrium solution to daisyworld can also be obtained by either integrating equation 5 or with a minimisation of x with respect to a_i subject to constraints. Each gives the same solution (Figs.4,5), independent of the history of the system. Thus *the equilibrium state in 2D and 0D daisyworld can be determined assuming the Gaia principle, without knowing detailed dynamics, just as the equilibrium states of thermodynamic systems can be determined by minimising free energy.*

VI. GAIA AND MEP IN THE LOGISTIC MAP

The logistic map is another widely used model for population growth (Hassell, Lawton and May, 1976). The resulting populations famously follow the period doubling route to chaos. There is no natural selection and again it is interesting to ask whether this model displays any kind of maximum population/maximum entropy behaviour.

The classical logistic map gives an n th population size of

$$y(n) = ay(n-1)[1 - y(n-1)] \quad (11)$$

The equilibrium distribution for y depends only on the parameter a . The mean population, averaged over 40000 iterations, is shown as a function of a in fig6. It exhibits a lot of structure which can be connected with the bifurcation diagram of the logistic map. The mean population of the attractor is higher than would be expected from maximum entropy (all levels equally occupied).

Turning now to entropy, there are many definitions related to the population distribution by analogy with the 2D daisyworld. For example, the Tsallis entropy (Latora et al 2000)

$$S = \sum_{i=1}^N \log(P(y_i)) / N \log(M/N) \quad (12)$$

where $P(y_i)$ is the number of occurrences of a population between $\frac{i-1}{N}$ and $\frac{i}{N}$ during M iterations of the logistic map (taken after a 1000 iteration equilibration period).

This quantity has a maximum possible value of 1, and (neglecting numerous distinct special cases of limit cycles) its value tends relatively smoothly to this value with increasing

a . However, its actual value depends on N . This only tells us that the area lying within the attractor increases with a .

The attractors in the logistic map provide some constraints on the mean value of y and the Tsallis entropy: it is difficult to define what the “maximum life” solution would be. Thus while it is notable that perturbed logistic map dynamics gives mean population in its stable region above 0.5 for all a , no quantitative principle emerges.

VII. CONCLUSIONS

The 2D-daisyworld model has been used as a testbed to determine whether any global maximisation is applicable to a system governed by the dynamics of natural selection. We find that Maximum Entropy Production is not valid here, either for biodiversity or heat transfer entropy. This casts doubt on some recent work in which the MEP is assumed as a constraint and used to obtain a solution (e.g. Pujol, 2003). Analysis of 2D and 0D cases suggests that daisyworlds appear to obey a separate optimisation principle, in the simple 2D case indistinguishable from MEP, that of maximising the amount of life present in the system. The logistic map attractors have above average population, but for this system it is difficult to define what the maximal population value should be.

The more fundamental implication is that stability of a dynamic equilibrium can be determined as an optimisation problem on the properties of the system as observed, and that detailed integration of system dynamics through time may be unnecessary in determining the equilibrium state.

VIII. REFERENCES

- P. Bak, C. Tang, and K. Wiesenfeld, "Self-organized criticality", Physical Review A 38 (1988) 364
- G. E. Crooks, "The Entropy Production Fluctuation Theorem and the Nonequilibrium Work Relation for Free Energy Differences", Phys. Rev. E 60, 2721 (1999),
- Dewar, R.C. Information theoretic explanation of maximum entropy production, the fluctuation theorem and self-organized criticality in non-equilibrium stationary states. J.Phys.A 36 631 (2003)

- Hassell, M. P., Lawton, J. H. and R. M. May. 1976. Patterns of dynamical behavior in single species populations. *J. Animal Ecology*. 45: 471-476.
- Kirchner, J.W. (2002). The Gaia hypothesis: Fact, theory, and wishful thinking. *Climatic Change* 52, 391-408.
- Lenton, T.M. (1998). Gaia and natural selection. *Nature* 394, 439-447.
- Lenton, T.M. and Lovelock, J.E. (2000). Daisyworld is Darwinian: Constraints on Adaptation are Important for Planetary Self-Regulation. *Journal of Theoretical Biology* 206, 109-114.
- Lorenz, R.D. (2002) Planets, life and the production of entropy. *International Journal of Astrobiology* 1, 3-13.
- Lorenz, R.D., Lunine, J.I., Withers, P.G. and McKay, C.P. (2001) Titan, Mars and Earth: Entropy Production by Latitudinal Heat Transport. *Geophys. Res. Lett.* 28, 415-418.
- Lovelock, J.E. (1988). *The Ages of Gaia - A Biography of Our Living Earth*. New York: W. W. Norton and Co.
- Latora, V., Baranger, M., Rapisarda, A. and C. Tsallis, C. "Generalization to nonextensive systems of the rate of entropy increase: the case of the logistic map", *Physics Letters A* 273 (2000) 97.
- May, R. M. 1976. Simple mathematical models with very complicated dynamics. *Nature*. 261: 459-467.
- Ozawa, H., Ohmura, A., Lorenz, R.D., Pujol, T. The second law of thermodynamics and the global climate system - A review of maximum entropy production principle. *Reviews of Geophysics* (accepted)
- Prasad, A., Mehra, V and Ramaswamy, R. Strange nonchaotic attractors in the quasiperiodically forced logistic map *Physical Review E*, 57, 1576-84 (1998)
- Pujol, T. (2002) The consequence of maximum thermodynamic efficiency in Daisyworld. *J. theor. Biol.* 217, 53-60.
- Robertson, D. and Robinson, J. (1998). Darwinian Daisyworld. *Journal of Theoretical Biology* 195, 129-134.
- Saunders, P.T. (1994). Evolution without Natural Selection: Further Implications of the Daisyworld Parable. *Journal of Theoretical Biology* 166, 365-373.
- Staley M Darwinian selection leads to Gaia in the long run *J Theor Biology* 218 35 2002
- Sugimoto, T. (2002). Darwinian evolution does not rule out Gaia hypothesis. *Journal of*

Theoretical Biology.218, 447-455.

Von Bloh, W., Block, A. and Schellnhuber, H.J. (1997) Self-stabilization of the biosphere under global change: a tutorial geophysiological approach. *Tellus* 49B, 249-262.

Von Bloh, W., Block, A. and Schellnhuber, H.J. (1999) Tutorial Modelling of geosphere-biosphere interactions: the effect of percolation-type habitat fragmentation. *Physica A* 266, 186-196.

Watson, A.J. and Lovelock, J.E. (1983). Biological homeostasis of the global environment: the parable of Daisyworld. *Tellus* 35B, 284-289.

Zipf, G.K Selective Studies and the Principle of Relative Frequency in Language (Harvard University Press, Cambridge, 1932)

-
- [1] This is implicit in using the sum over all paths to build a partition function
 - [2] Finite diffusivity leads to clump formation, which confounds this result due to neglect the entropy associated with variation in clump size. In principle this could be incorporated, but for the present purposes the maximum entropy principle observed in the limit of infinite diffusivity is sufficient.
 - [3] We say "close to": these are the limiting cases for low death rates and high birth rates, assuming uncorrelated dead patches.
 - [4] factor of two arises from double counting




FIG. 1: Fractional daisy coverage and temperature (in units of optimal growth temperature, $T_g = 295.5$) from a calculation with a 200×200 lattice ($\gamma = 0.02(1 - (\delta + T_d - T)(T + \delta - T_d)/2\delta^2$, $\beta(T) = (\delta + T_g - T)(T + \delta - T_g)/\delta^2$. $T_d = 305.5$, $\delta = 17.5$) $D = 0.1$, $r = 0.02$) with scaled insolation S varied from 0.7 to 1.7 over 10^6 updates of daisy and temperature fields with a temperature dependent death rate, minimal at $T_d = 305.5$. The graph shows that in the non-desert regime population tends to its maximal value ($1 - x = \gamma/(\gamma + \beta) = 0.99$ in the mean field approximation, where regrowth is always possible) while temperature is regulated at T_d , rather than T_g which would maximise(bio-)entropy production. Thus the governing global principal appears to be Gaian (maximal life) rather than MEP. At higher S , fluctuations lead to long lived deserts, which cause the mean field approximation to break down as daisies cannot grow in the interior.




FIG. 2: Graph showing net total energy flows into (solid line) and out of (dotted line) a 200x200 cell 2D daisyworld system (diffusion = 0.1, death = 0.02, mutation = 0.02). The simulation starts with $S=1.0$ and increases by discrete jumps of 0.1 every 10000 timesteps. Up to $S=1.7$ this is sufficient to allow the system to return to equilibrium: the excess incoming flow of energy is reduced by evolution of the albedo. Beyond 1.7 (see inset) the sudden change generates sufficient die back to create deserts in the system, which initially cause positive feedback and increase the incoming flow still further. Ultimately, evolution of paler daisies in the non-desert regions allow reinvansion of the deserts, but the simulation had to be run for 20000 timesteps ($S=1.7$) and 50000 ($S=1.8$) to achieve this. Even in equilibrium, transient desert regions occur. At $S=1.9$ the simulation was run for 120000 timesteps, throughout which desert and populated regions coexisted and migrated around the planet (there being no curvature to favour one region over another). Throughout, note that the energy emitted is smoother than that absorbed and lags the energy absorbed by about 500 timesteps (10 mean daisy lifetimes) $\gamma = 0.02, D = 0.1, r = 0.02$.




FIG. 3: Graphs showing variation of configuration (biodiversity) entropy $\langle \sum_{i=1}^{20} p(A_i) \ln p(A_i) \rangle$ from partitioning the albedos into 20 groups, and (heat) entropy production $\langle \int dQ/T \rangle$ averaged over the 200x200 lattice and over a period of 50 timesteps (one mean daisy lifetime) for the same calculation as Fig.2. Each changes monotonically as the temperature is increased through the region of full coverage. Beyond $S=1.7$, where fluctuating deserts appear, each measure becomes less well defined. There is significantly increased heat flow from the hot deserts to the covered regions, so the heat flow becomes dominated by the area of desert. Conversely, the biodiversity is reduced when much of the planet is unpopulated. Reversing the change in temperature shows that the entropy change is reversible.




FIG. 4: Results from 0D daisyworld with parameter values from Lenton and Lovelock: $S/\sigma_B = 1.68 \times 10^{10} K^4$, $q = 2.06425 \times 10^9$ and $\gamma = 0.3$. a) Temperature variation with increasing solar insolation. Thick line show planetary average, dotted lines show T_b and T_w . The four regimes are shown in their region of stability: Simulations commencing with temperatures outside the growing range can stabilise the bare planet solution, but single-daisy and two-daisy solutions do not coexist. Only stable solutions are shown, including the bistable living/dead regime. b) coverage: Thick line denotes bare ground, thin lines denote individual species. For non-zero a_b and a_w , $da_i/dt = 0$ means that $\beta_b = \beta_w$ and from 7 and 9, T_w and T_b are also independent of L : $T_w = 22.5$ for $q = 0$ falling almost linearly to zero for $q = 9.34 \times 10^9$. For standard parameter values from Watson and Lovelock, $\beta_b = \beta_w = 0.918$ $T_w=17.5$ and $T_b=27.5$. With T_w and β_w fixed by the choice of q , independent of L , as is the amount of bare ground at equilibrium: $x = \gamma/\beta_w$; (from 1). Equations 6 and 8 now give us simple linear expressions for a_b and a_w in terms of x and the single variable A_p . Thus, once the assumption of species coexistence is made, the only L -dependent variables are




FIG. 5: Daisyworld with additional grey daisy species with albedo 0.5. Other parameters as for Fig 4. a) Temperature - thick line shows planetary average, thin lines denote local temperatures over daisies. Note that the grey-only solution is stable because it maximises life, despite having no temperature regulatory effect. b) Coverage: Thick line denotes bare ground, thin lines denote individual species. Note that graph is continuous, with bare ground (x) falling below 0.30625 only in the single-species regions.




FIG. 6: Bold line shows the mean amount of life from the logistic map, plotted against recursion parameter a . Background dots show the bifurcation diagram. A small amount of random noise added to $y(n)$ (drawn from a flat distribution ± 0.001) eliminates most of the ordered behaviour, and the gives mean population above 0.5 throughout. Without noise, the mean population is still above 0.5 for all the chaotic cases, and for all the ordered windows except for the tiny region between 3.9602 and 3.9616, 0.04% of the total. Populations above 0.5 are found for stable attractors across the family of logistic maps of the form $y_{n+1} = a[y_n(1 - y_n)]^\nu$